arXiv:1301.0450v1 [math.CO] 3 Jan 2013

Long properly colored cycles in edge-colored complete graphs *

Guanghui Wang[a], Tao Wang[b, c], Guizhen Liu[a]

[a]School of Mathematics,
Shandong University, 250100, Jinan, Shandong, P. R. China
[b]Institute of Applied Mathematics
Henan University, Kaifeng, 475004, P. R. China
[c]College of Mathematics and Information Science
Henan University, Kaifeng, 475004, P. R. China

Abstract

Let K_n^c denote a complete graph on n vertices whose edges are colored in an arbitrary way. Let $\Delta^{\mathrm{mon}}(K_n^c)$ denote the maximum number of edges of the same color incident with a vertex of K_n. A properly colored cycle (path) in K_n^c is a cycle (path) in which adjacent edges have distinct colors. B. Bollobás and P. Erdös (1976) proposed the following conjecture: If $\Delta^{\mathrm{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, then K_n^c contains a properly colored Hamiltonian cycle. Li, Wang and Zhou proved that if $\Delta^{\mathrm{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, then K_n^c contains a properly colored cycle of length at least $\lceil \frac{n+2}{3} \rceil + 1$. In this paper, we improve the bound to $\lceil \frac{n}{2} \rceil + 2$.

1 Introduction

All graphs considered here are finite, undirected, and simple. Let G be a graph with vertex set V and edge set E.

An *edge coloring* of a graph is an assignment of "colors" to the edges of the graph. An *edge colored graph* is a graph which is assigned a color for each edge. A cycle (path) in an edge colored graph is *properly colored* if no two incident edges in it have the same color.

Grossman and Häggkvist [9] gave a sufficient condition on the existence of a properly colored cycles in edge colored graphs with two colors, and Yeo [16] extended the result to edge colored graphs with any number of colors. Given an edge-colored graph G, let $\deg^c(v)$, named the color degree of a vertex v, be defined as the maximum number of edges adjacent to v, that have distinct colors. The minimum color degree $\delta^c(G)$ is the minimum $\deg^c(v)$ over all vertices of G. In [12], some minimum color degree

---

*This research was supported by NSFC Grants (61070230, 11101243, 10901097, 11101125) and RFDP(20100131120017).

conditions for the existence of properly colored cycles are obtained. In particular, they proved that if G is an edge colored graph of order n satisfying $\delta^c(G) \geq d \geq 2$, then either G has a properly colored path of length at least $2d$, or G has a properly colored cycle of length at least $\lceil \frac{2d}{3} \rceil + 1$. In [13], Lo improved the bound $\lceil \frac{2d}{3} \rceil + 1$ to the best possible value $d + 1$.

Let K_n^c be an edge colored complete graph on n vertices, the color assigned to edge uv is $c(u, v)$. In [4], Barr gave a simple sufficient condition for the existence of a properly colored Hamiltonian path in K_n^c: K_n^c has no monochromatic triangles. Bang-Jensen, Gutin and Yeo [3] proved that if K_n^c contains a properly colored 2-factor, then it has a properly colored Hamiltonian path. In [7], Feng et al. proved that K_n^c has a properly colored Hamilton path iff K_n^c has a properly colored almost 2-factor (an almost 2-factor is a spanning subgraph consisting of disjoint cycles and a path).

Bollobás and Erdös [5] proved that if $n \geq 3$ and $\delta^c(K_n^c) \geq \frac{7n}{8}$, then there exists a properly colored Hamiltonian cycle. They also proposed a question: whether the bound $\frac{7n}{8}$ can be improved to $\frac{n+5}{3}$. Fujita and Magnant [8] construct an edge colored complete graph K_{2m} with $\delta^c(K_{2m}) = m$, which has no properly colored Hamiltonian cycles. So $\delta^c(K_n) = \lfloor \frac{n}{2} \rfloor$ is not enough. Let $\Delta^{\text{mon}}(K_n^c)$ be the maximum number of edges of the same color incident with a vertex of K_n^c, an edge colored complete graph on n vertices. In the same paper, Bollobás and Erdös proposed the following conjecture.

Conjecture 1 (Bollobás and Erdös [5]). If $\Delta^{\text{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, then K_n^c has a properly colored Hamiltonian cycle.

Bollobás and Erdös proved that if $\Delta^{\text{mon}}(K_n^c) \leq \frac{n}{69}$ then K_n^c contains a properly colored Hamiltonian cycle. This result was improved by Chen and Daykin [6] and Shearer [15].

As far as we know, the best asymptotic estimate was obtained by Alon and Gutin using the probabilistic method.

Theorem 1.1 (Alon and Gutin [1]). For every positive real number ϵ, there exists an $n_0 = n_0(\epsilon)$ so that for every $n > n_0$, if K_n^c satisfies

$$\Delta^{\text{mon}}(K_n^c) \leq (1 - \frac{1}{\sqrt{2}} - \epsilon)n, \tag{1.1}$$

then K_n^c contains a properly colored Hamiltonian cycle.

Li, Wang and Zhou [12] studied long properly colored cycles in edge colored complete graphs and proved that if $\Delta^{\text{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, then K_n^c contains a properly colored cycle of length at least $\lceil \frac{n+2}{3} \rceil + 1$. For more details concerning properly colored cycles and paths, we refer the reader to the survey papers [2, 10, 11]. In this paper, we improve the bound on the length of the properly colored cycles and prove the following theorem.

Theorem 1.2. If $\Delta^{\text{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, then K_n^c contains a properly colored cycle of length at least $\lceil \frac{n}{2} \rceil + 2$.

The main idea is following the rotation-extension technique of Pósa [14], which was used on general edge colored graphs in [13].

When a and b are integers, the notation $[a, b]$ is used to indicate the interval of all integers between a and b, including both. In particular, $[a, a-1]$ is an empty set. A properly colored path P of length ℓ is viewed as an ℓ-tuple $(v_0, v_1, \ldots, v_\ell)$, which is different from $(v_\ell, v_{\ell-1}, \ldots, v_0)$. For such a properly colored path $P = (v_1, v_2, \ldots, v_\ell)$, denote $\{v \mid c(v_1, v_2) \neq c(v_1, v)\}$ by $N^c(v_1, P)$ and $\{v \mid c(v_\ell, v_{\ell-1}) \neq c(v, v_\ell)\}$ by $N^c(v_\ell, P)$.

2 The Proof of Theorem 1.2

If $3 \leq n \leq 5$, then K_n^c is a properly colored complete graph, and hence each Hamiltonian cycle is properly colored. Our conclusion holds clearly. So we may assume that $n \geq 6$.

We will prove Theorem 1.2 by contradiction. Suppose that each properly colored cycle is of length at most $\lceil \frac{n}{2} \rceil + 1$. For simplicity, let the vertices of K_n^c be labeled with integers from 1 to n. Let P_0 be a longest properly colored path. Without loss of generality, assume that $P_0 = (1, 2, \ldots, \ell)$.

Put

$$X = \{x_1, x_2, \ldots, x_p\} = \{x_i \mid c(1, x_i) \neq c(1, 2)\} = N^c(1, P_0)$$

and

$$Y = \{y_1, y_2, \ldots, y_q\} = \{y_i \mid c(y_i, \ell) \neq c(\ell - 1, \ell)\} = N^c(\ell, P_0).$$

Moreover, $x_1, x_2, \ldots, x_p$ and $y_1, y_2, \ldots, y_q$ are increasing sequences. Since $\Delta^{\mathrm{mon}}(K_n^c) < \lfloor \frac{n}{2} \rfloor$, we have $\min\{|X|, |Y|\} \geq \lceil \frac{n}{2} \rceil$, consequently, $x_1 < x_p$. Note that $X, Y \subset V(P_0)$ since P_0 is a longest properly colored path. Thus $\ell \geq \lceil \frac{n}{2} \rceil + 2$. Note that either $\ell \notin X$ or $1 \notin Y$, otherwise $(1, 2, \ldots, \ell, 1)$ is a properly colored cycle of length at least $\lceil \frac{n}{2} \rceil + 2$, which is a contradiction. Hence $\ell \geq \lceil \frac{n}{2} \rceil + 3$.

Let $y_s \in Y$ be the maximum such that $c(\ell, y_i) = c(y_i, y_i + 1)$ for all $y_i \in [y_1, y_s] \cap Y$. Note that y_s is well defined, since $c(\ell, y_1) = c(y_1, y_1 + 1)$, otherwise, $(y_1, y_1 + 1, \ldots, \ell, y_1)$ is a properly colored cycle of length at least $\lceil \frac{n}{2} \rceil + 2$. Clearly, $y_s \leq \ell - 2$.

Note that $c(1, x_p) = c(x_p - 1, x_p)$.

Claim 1. $y_s \leq x_p - 3$.

Proof of Claim 1. If $x_p = \ell$, then $y_s \leq \ell - 3$, otherwise, $(1, 2, \ldots, \ell - 2, \ell, 1)$ is properly colored cycle of length $\ell - 1 \geq \lceil \frac{n}{2} \rceil + 2$, a contradiction. So we may assume that $x_p \leq \ell - 1$. Let $y_i \in Y$ be the maximum such that $y_i < x_p$. If $c(y_i, y_i + 1) = c(y_i, \ell)$, then $(1, \ldots, y_i, \ell, \ell - 1, \ldots, x_p, 1)$ is a properly colored cycle containing $Y \cup \{\ell, \ell - 1\}$, a contradiction. Hence, $c(y_i, y_i + 1) \neq c(y_i, \ell)$, by the definition of y_s, we have $y_s \leq x_p - 2$. If $y_s = x_p - 2$, then $(1, \ldots, x_p - 2, \ell, \ell - 1, \ldots, x_p, 1)$ is a properly colored cycle of length $\ell - 1 \geq \lceil \frac{n}{2} \rceil + 2$, which is a contradiction. Therefore, $y_s \leq x_p - 3$. □

Claim 2. There exist $u, w \in X$ such that $1 \leq y_1 \leq y_s < u \leq \lceil \frac{n}{2} \rceil + 1$, $u < w$ and the following proposition holds:

(a) $c(y_i, \ell) = c(y_i, y_i + 1)$ for all $y_i \in [y_1, y_s] \cap Y$;

(b) $c(1, x_i) = c(x_i, x_i + 1)$ for all $x_i \in [y_1 + 1, u] \cap X$;

(c) $c(1, w) \neq c(w, w+1)$ or $w = \ell$;

(d) if $x_i \in X$ and $x_i < w$, then $x_i \leq u$.

Proof of Claim 2. (a) By the definition of y_s, the assertion (a) holds.

(b) First, we show that $c(1, x_i) = c(x_i, x_i+1)$ for all $x_i \in [y_1+1, y_s+1] \cap X$. Otherwise, if there exists $x_i \in [y_1+1, y_s+1] \cap X$ such that $c(1, x_i) \neq c(x_i, x_i+1)$, let y_j be the maximum such that $y_j \in [y_1, y_s] \cap Y$ and $y_j < x_i$, then $(1, 2, \ldots, y_j, \ell, \ell-1, \ldots, x_i, 1)$ is a properly colored cycle containing $Y \cup \{\ell-1, \ell\}$, a contradiction.

Next, we show that there exists u such that $c(1, x_i) = c(x_i, x_i+1)$ for all $x_i \in [y_s+1, u] \cap X$. Let $x \in X$ be the minimum such that $x > y_s$. Note that x must be exists since $x_p > y_s$ by Claim 1. If $x = \ell$, then $y_s \geq y_1 \geq 2$, hence $(1, \ldots, y_s, \ell, 1)$ is a properly colored cycle containing $X \cup \{1, 2\}$, a contradiction, so $x < \ell$. Suppose that $c(1, x) \neq c(x, x+1)$. If $y_s = 1$, then $(1, \ell, \ell-1, \ldots, x, 1)$ is a properly colored cycle containing $X \cup \{1, \ell\}$, a contradiction; if $y_s > 1$, then $(1, \ldots, y_s, \ell, \ell-1, \ldots, x, 1)$ is a properly colored cycle containing $X \cup \{1, 2\}$, a contradiction. So, we have $x < \ell$ and $c(1, x) = c(x, x+1)$.

Let u be the maximum such that $c(1, x_i) = c(x_i, x_i+1)$ for all $x_i \in [y_s+1, u] \cap X$ and $y_s < u < \ell$. By the above argument, u is well defined. In particular, $c(1, u) = c(u, u+1)$, it follows that $(1, 2, \ldots, u, 1)$ is a properly colored cycle, and hence $u \leq \lceil \frac{n}{2} \rceil + 1$.

Let $w \in X$ be the minimum such that $w > u$. Clearly (c) and (d) hold. □

Let $S(P_0) = Y \cap [y_1, y_s]$. Note that y_1, y_s, u, w, S are viewed as functions of longest properly colored paths.

By Claims 2 (a) and 2 (c), $C_0 = (1, \ldots, y_s, \ell, \ell-1, \ldots, w, 1)$ is a properly colored cycle, see Fig. 1.



Fig. 1: Cycles $C_0 = (1, \ldots, y_s, \ell, \ell-1, \ldots, w, 1)$

Without loss of generality, assume that P_0 is a longest properly colored path satisfying $|S(P_0)|$ is maximum over all the longest properly colored paths.

Clearly, $P^* = (y_1+1, y_1+2, \ldots, \ell, y_1, \ldots, 1)$ is a longest properly colored path. Note that

$$N^c(1, P^*) = X.$$

Since $c(y_1+1, y_1+2) \neq c(y_1+1, y_1)$, it follows that $y_1 \in N^c(y_1+1, P^*)$. Hence P^* is a longest properly colored path.

Claim 3. $|X| = \lceil \frac{n}{2} \rceil$ and $|C_0| = \lceil \frac{n}{2} \rceil + 1$. Moreover, $S(P_0) = [y_1, y_s]$, i.e., it is an integer interval; and

$$X = \begin{cases} [3, y_1] \cup [t, u] \cup [w, \ell] & \text{if } y_1 > 1; \quad (2.1) \\ [t, u] \cup [w, \ell-1] & \text{if } y_1 = 1, \quad (2.2) \end{cases}$$

where $|[t, u]| = |[y_1, y_s]|$, $t \geq \max\{y_1+1, 3\}$ and $S(P^*) = [t, u]$.

Proof of Claim 3. By Claim 2 (b), $[y_1+1,u]\cap X\subseteq S(P^*)$. Since $c(1,u)=c(u,u+1)$, it yields that $u\in S(P^*)$. If $w<\ell$, then $c(1,w)\neq c(w,w+1)$, thus $w\notin S(P^*)$; if $w=\ell$, then $c(1,\ell)\neq c(\ell,y_1)$, thus $w\notin S(P^*)$. Therefore, $S(P^*)=[y_1+1,u]\cap X$ and $|[y_1,y_s]\cap Y|\geq|[y_1+1,u]\cap X|$ by the maximality of $|S(P_0)|$.

Now

$$\begin{aligned}
\left\lceil\frac{n}{2}\right\rceil+1 &\geq |C_0| \\
&= |[1,y_s]\cap X|+|[w,\ell]\cap X|+|[1,y_s]\setminus X|+|[w,\ell]\setminus X| \\
&= |X|-|[y_s+1,u]\cap X|+|[1,y_s]\setminus X|+|[w,\ell]\setminus X| \\
&= |X|-|[y_1+1,u]\cap X|+|[y_1+1,y_s]\cap X|+|[1,y_s]\setminus X|+|[w,\ell]\setminus X| \\
&\geq |X|-|[y_1,y_s]\cap Y|+|[y_1+1,y_s]|+|[1,y_1]\setminus X|+|[w,\ell]\setminus X| \\
&= |X|-|[y_1,y_s]\cap Y|+|[y_1,y_s]|+|[2,y_1]\setminus X|+|[w,\ell]\setminus X| \\
&\geq |X|+|[2,y_1]\setminus X|+|[w,\ell]\setminus X| \\
&\geq |X|+1 \\
&\geq \left\lceil\frac{n}{2}\right\rceil+1.
\end{aligned}$$

Therefore, all the inequalities become equalities. Then $|C_0|=\lceil\frac{n}{2}\rceil+1$, $|X|=\lceil\frac{n}{2}\rceil$, $|S(P_0)|=|S(P^*)|$, $S(P_0)=[y_1,y_s]$ and

$$|[2,y_1]\setminus X|+|[w,\ell]\setminus X|=1. \tag{2.3}$$

(2.3) implies that

$$\begin{cases} [3,y_1]\subseteq X \text{ and } [w,\ell]\subseteq X & \text{if } y_1>1; \qquad (2.4)\\ [w,\ell]\cap X=[w,\ell-1] & \text{if } y_1=1. \qquad (2.5)\end{cases}$$

Since $|S(P_0)|=|S(P^*)|$, by the above argument, $S(P^*)$ must be an integer interval, so $S(P^*)=[y_1+1,u]\cap X=[t,u]$, clearly $t\geq\max\{y_1+1,3\}$. □

Note that t is also viewed as a function of the longest properly colored path.

Claim 4.

(a) For any $y\in Y\cap[y_s+1,w-1]$, we have $c(y,y-1)=c(y,\ell)$; $|S(P_0)|\geq|Y\cap[y_s+1,w-1]|$.

(b) If $w=x_p$, then $X=[3,\lceil\frac{n}{2}\rceil+1]\cup\{x_p\}$.

Proof of Claim 4. (a) Since $|C_0|=\lceil\frac{n}{2}\rceil+1$, we have $c(y,y-1)=c(y,\ell)$ for any $y\in Y\cap[y_s+1,w-1]$; otherwise, $(1,2,\dots,y,\ell,\ell-1,\dots,w,1)$ is a properly colored cycle of length greater than $|C_0|$, a contradiction.

Obviously, $Q=(w-1,w-2,\dots,1,w,w+1,\dots,\ell)$ is a longest properly colored path. Note that $N^c(\ell,Q)=Y$. We have just proved $Y\cap[y_s+1,w-1]\subseteq S(Q)$, and hence $|S(P_0)|\geq|S(Q)|\geq Y\cap[y_s+1,w-1]$.

(b) Since $w=x_p$, it follows that $u=x_{p-1}$ and $(1,2,\dots,x_{p-1})$ is a properly colored cycle containing $\{1,2,x_1,\dots,x_{p-1}\}$, so $x_{p-1}=\lceil\frac{n}{2}\rceil+1$. Hence $X=[3,\lceil\frac{n}{2}\rceil+1]\cup\{x_p\}$. □

Claim 5. $|S(P_0)| \geq 3$.

Proof of Claim 5. By way of contradiction, we assume that $|S(P_0)| \leq 2$. Without loss of generality, we may assume that $y_1 \neq 1$, otherwise consider the properly colored path $(2, 3, \ldots, \ell, 1)$ instead.

Case 1: $w \leq \ell - 1$.

Let $P' = (\ell, \ell - 1, \ldots, 1)$, it is a longest properly colored path. If $w \leq \ell - 1$, then $c(1, \ell) = c(\ell, \ell - 1)$ and $c(1, \ell - 1) = c(\ell - 1, \ell - 2)$, so $\ell, \ell - 1 \in S(P')$. Hence, $|S(P_0)| = |S(P')| = 2$ and $c(1, \ell - 2) \neq c(\ell - 2, \ell - 3)$. Applying Claim 3 with P', we have $c(\ell, \ell - 1) \neq c(\ell, 2)$ and $y_1 = 2$, and hence $y_s = 3$.

Case 1.1: $w \leq \ell - 2$. Then $C_1 = (1, 2, \ldots, \ell - 2, 1)$ is a properly colored cycle of length at most $\lceil \frac{n}{2} \rceil + 1$, i.e., $\ell - 2 \leq \lceil \frac{n}{2} \rceil + 1$, or $\ell \leq \lceil \frac{n}{2} \rceil + 3$. On the other hand, $\ell \geq \lceil \frac{n}{2} \rceil + 3$, then $\ell = \lceil \frac{n}{2} \rceil + 3$. Moreover, $Y = [2, \ell - 2]$. Since the properly colored cycle $C_0 = (1, 2, 3, \ell, \ell - 1, \ldots, w, 1)$ has length $\lceil \frac{n}{2} \rceil + 1$ by Claim 3, we conclude that $w = 6$. Note that $|X| = \lceil \frac{n}{2} \rceil = |[t, u]| + |[w, \ell]| \geq 5$ implying $n \geq 9$.

If $\ell = n$, then $n - 2 = \ell - 2 = \lceil \frac{n}{2} \rceil + 1$ and $n = 6, 7$, a contradiction. Then we may assume that there exists a vertex z such that it is not on P_0. The properly colored path $(1, 2, 3, \ell, \ell - 1, \ldots, 5, 4)$ is longest, it follows that $c(4, 5) = c(4, z)$, otherwise, $(1, 2, 3, \ell, \ell - 1, \ldots, 5, 4, z)$ is a longer properly colored path. By Claim 4(a), we have $c(\ell, 5) = c(5, 4) \neq c(5, 6)$. Therefore, $(\ell - 1, \ell, 5, 6, \ldots, \ell - 2, 1, 2, 3, 4, z)$ is a longer properly colored path than P_0, which is a contradiction.

Case 1.2: $w = \ell - 1$. Then $|X| = \lceil \frac{n}{2} \rceil = |[t, u]| + |[w, \ell]| = 4$, i.e., $n = 7, 8$. Since $\left\lceil \frac{n}{2} \right\rceil + 3 \leq \ell \leq n$, $\ell = 7$ or $\ell = 8$.

If $t = 3$, then $c(1, 3) = c(3, 4)$. Since $y_s = 3$, $c(3, 4) = c(\ell, 3)$. Consider the properly colored path $P'' = (3, 4, \ldots, \ell, 2, 1)$. By Claim 3, we have $S(P'') = \{3, 4\}$ and $N^c(3, P'') = \{t', t' + 1\} \cup \{\ell, 2\}$, so $c(\ell, 3) \neq c(3, 4)$, a contradiction.

If $t = 4$, then $X = \{4, 5, \ell - 1, \ell\}$. Define ϕ be the permutation on $[1, \ell]$ such that $(\phi(1), \phi(2), \ldots, \phi(\ell)) = (3, 4, \ldots, \ell, 2, 1)$. We will show the following propositions hold for $i \geq 0$ by induction on i.

(a) $P_i = (\phi^i(1), \phi^i(2), \ldots, \phi^i(\ell))$ is a longest properly colored path;

(b) $S(P_i) = \{\phi^i(2), \phi^i(3)\}$;

(c) $N^c(\phi^i(1), P_i) = \{\phi^i(j) \mid j \in [4, 5] \cup [\ell - 1, \ell]\}$.

(d) $c(\phi^i(1), \phi^i(j)) = c(\phi^i(j), \phi^i(j + 1))$ for $j \in [4, 5]$.

Clearly, the statements (a)–(d) hold for $i = 0$. Assume that $i \geq 1$ and they are true for $i - 1$. Let j' denote $\phi^{i-1}(j)$. Since $2' \in S(P_{i-1})$, $c(1', 2') \neq c(2', 3') = c(2', \ell')$. Hence, $P_i = (\phi(1'), \phi(2'), \cdots, \phi(\ell'))$ is a longest properly colored path. Also, by the form of $N^c(\phi^{i-1}(1), P_{i-1})$ and $|S(P_i)| = |S(P_0)| = 2$, (b) holds. By Claim 3 and $|X| = 4$,

$$N^c(\phi^i(1), P_i) = \{\phi^i(j) \mid j \in [t_i, t_i + 1] \cup [\ell - 1, \ell]\},$$

for some $t_i = 3, 4$. If $t_i = 3$, we can also get a contradiction as above by taking $P_0 = P_i$. So $t_i = 4$, and thus (c) holds. By Claim 3, (d) also holds.

If $\ell = 7$, then $X = \{4, 5, 6, 7\}$ and $Y = \{2, 3, 4, 5\}$. For the path P_0, we have $c(7, 2) = c(2, 3)$ and $c(7, 3) = c(3, 4)$. Taking $i = 1$ in the statement, consider the longest properly colored path P_1, i.e., $(3, 4, 5, 6, 7, 2, 1)$. By statement (c), $c(\phi^1(1), \phi^1(5)) = c(\phi^1(5), \phi^1(6))$, i.e., $c(3, 7) = c(7, 2)$. Therefore, $c(2, 3) = c(7, 2) = c(3, 7) = c(3, 4)$, a contradiction.

If $\ell = 8$, then $X = \{4, 5, 7, 8\}$. Since $u = 5$, it follows that $c(1, 5) = c(5, 6) \neq c(4, 5)$. Consider the properly colored path $P_5 = (4, 3, 6, 5, 8, 7, 1, 2)$. Since $8 = \phi^5(5, P_5)$, by statement (c) and Claim 2 (b), $c(4, 8) = c(8, 7) \neq c(5, 8)$; moreover, $5 \in N^c(8, P_0)$. Thus $c(8, 5) = c(4, 5)$ by Claim 4. Hence $c(1, 5) = c(5, 6) \neq c(4, 5) = c(5, 8)$. Now, we have a properly colored cycle $(1, 2, 3, 4, 8, 5, 1)$ of length 6, a contradiction.

Case 2: $w = \ell$. Since $|Y \cap [y_s + 1, w - 1]| \leq |S(P_0)|$ by Claim 4 (a), it follows that $|Y| = |[y_1, y_s]| + |Y \cap [y_s + 1, w - 1]| \leq 2|S(P_0)|$. If $|S(P_0)| = 1$, then $\lceil \frac{n}{2} \rceil \leq |Y| \leq 2$, and thus $n \leq 4$, a contradiction. Hence, $|S(P_0)| = 2$, and $\lceil \frac{n}{2} \rceil \leq |Y| \leq 4$, so $n \leq 8$. By Claim 3, $X = [3, \lceil \frac{n}{2} \rceil + 1] \cup \{\ell\}$, $u = \lceil \frac{n}{2} \rceil + 1$, $t = \lceil \frac{n}{2} \rceil$, $y_1 = t - 1 = \lceil \frac{n}{2} \rceil - 1$. So $Y \subseteq [\lceil \frac{n}{2} \rceil - 1, \ell - 2]$, hence, $\lfloor \frac{n}{2} \rfloor = n - 2 - (\lceil \frac{n}{2} \rceil - 1) + 1 \geq \ell - 2 - (\lceil \frac{n}{2} \rceil - 1) + 1 \geq |Y| \geq \lceil \frac{n}{2} \rceil$, then $n = \ell$ is even, $Y = [\frac{n}{2} - 1, n - 2]$ and $|Y| = \frac{n}{2}$. Now, we know that $n = 6, 8$.

If $n = 6$, then $X = \{3, 4, 6\}$, $Y = \{2, 3, 4\}$, $y_1 = 2$, $y_s = t = 3$ and $u = 4$. By Claim 2 (a) and 2 (b), $c(1, 3) = c(3, 4) = c(3, 6)$, which contradicts the fact that $\Delta^{\text{mon}}(K_6) < 3$.

If $n = 8$, then $X = \{3, 4, 5, 8\}$, $Y = \{3, 4, 5, 6\}$, $y_1 = 3$, $t = 4$ and $u = 5$. Since $4 \in [y_1 + 1, u] \cap X$, $c(1, 4) = c(4, 5) \neq c(3, 4)$ by Claim 2 (b); by Claim 2 (a), $c(8, 3) = c(3, 4) \neq c(2, 3)$ and $c(8, 4) = c(4, 5) \neq c(3, 4)$. Hence, $c(4, 1) = c(4, 5) = c(4, 8)$. Since $\Delta^{\text{mon}}(K_8) < 4$, it follows that $c(4, 7) \neq c(4, 1)$. If $c(4, 7) \neq c(7, 8)$, then $(1, 2, 3, 8, 7, 4, 1)$ is a properly colored cycle of length 6, a contradiction. So $c(4, 7) = c(7, 8) \neq c(7, 6)$. By Claim 4 (a), we have $c(8, 5) = c(4, 5) \neq c(5, 6)$, so $c(8, 5) \neq c(8, 3)$. Now we have a properly colored Hamiltonian cycle $(1, 2, 3, 8, 5, 6, 7, 4, 1)$, a contradiction. □

Claim 6. Without loss of generality, we may assume that $t \geq y_1 + 3$.

Proof of the Claim 6. Suppose to the contrary that $t \leq y_1 + 2$. Then $t = y_1 + 1$ or $t = y_1 + 2$. Since $|S(P_0)| \geq 3$, we have $t - 1 \in S(P_0)$. Without loss of generality, we may assume that $y_1 = 1$, otherwise, consider the longest properly colored path $(t, t + 1, \ldots, \ell, t - 1, t - 2, \ldots, 1)$ instead. By Claim 3, $t = 3$. Moreover, $X = [3, y_s + 2] \cup [w, \ell - 1]$.

Let ϕ be a permutation on $[1, \ell]$ such that $(\phi(1), \phi(2), \ldots, \phi(\ell - 2), \phi(\ell - 1), \phi(\ell)) = (3, 4, \ldots, \ell, 2, 1)$. We will show the following statements hold for $i \geq 0$ by induction.

(a) $P_i = (\phi^i(1), \phi^i(2), \ldots, \phi^i(\ell))$ is a longest properly colored path;

(b) $S(P_i) = \{\phi^i(j) \mid j \in [1, y_s]\}$;

(c) $N^c(\phi^i(1), P_i) = \{\phi^i(j) \mid j \in [3, y_s + 2] \cup [w, \ell - 1]\}$;

(d) for all $j \in [3, y_s + 2]$, we have $c(\phi^i(1), \phi^i(j)) = c(\phi^i(j), \phi^i(j + 1))$;

(e) for all $j \in [4, \lceil \frac{n}{2} \rceil + 2]$, we have $c(\phi^i(2), \phi^i(j)) = c(\phi^i(j), \phi^i(j + 1))$.

First, the statements are true for $i = 0$. Assume that the statements are true for $i-1$, where $i \geq 1$. Since $\phi^{i-1}(2) \in S(P_{i-1})$, it follows that $P_i = (\phi^{i-1}(3), \phi^{i-1}(4), \ldots, \phi^{i-1}(\ell), \phi^{i-1}(2), \phi^{i-1}(1)) = (\phi^i(1), \phi^i(2), \ldots, \phi^i(\ell))$ is a longest properly colored path. Since $|S(P_{i-1})| = |S(P_0)| = |[1, y_s]|$, we have $t(P_{i-1}) = \phi^{i-1}(3)$; by Claim 3, it yields that $S(P_i) = [t(P_{i-1}), u(P_{i-1})] = [\phi^i(1), \phi^i(y_s)]$, and thus $|S(P_i)| = |S(P_0)|$. Hence, by the assumption, we have $N^c(\phi^i(1), P_i) = \{\phi^i(j) \mid j \in [3, y_s+2] \cup [w, \ell-1]\}$. By Claim 2 (b), for all $j \in [3, y_s+2]$, $c(\phi^i(1), \phi^i(j)) = c(\phi^i(j), \phi^i(j+1))$.

Since $S(P_i) = \{\phi^i(j) \mid j \in [1, y_s]\}$, it follows that $P_i' = (\phi^i(2), \phi^i(3), \ldots, \phi^i(\ell), \phi^i(1))$ is a longest properly colored path. By Claim 3, $S(P_i') = \{\phi^i(j) \mid j \in [3, y_s+2]\}$. By Claim 2 (a), $c(\phi^i(2), \phi^i(3)) = c(\phi^i(2), \phi^i(\ell))$, so $\phi^i(\ell) \notin N^c(\phi^i(2), P_i')$; by Claim 3, $N^c(\phi^i(2), P_i') = \{\phi^i(j) \mid j \in [4, \lceil \frac{n}{2} \rceil + 2] \cup \{1\}\}$. By Claim 2 (b), $c(\phi^i(2), \phi^i(j)) = c(\phi^i(j), \phi^i(j+1))$ for all $j \in [4, \lceil \frac{n}{2} \rceil + 2]$. This completes the proof of the statements.

Consider the properly colored path P_0. Since $|S(P_0)| \geq 3$, it follows that $y_s \geq 3$. Thus $5 \in [3, y_s+2]$. By statement (d), $c(1,5) = c(5,6)$.

If ℓ is odd, take $i = \frac{\ell+1}{2}$. It is easy to check that $P_{\frac{\ell+1}{2}} = (1,4,3,6,5,\ldots)$. By statement (d), $c(1,5) \neq c(6,5)$, a contradiction.

If ℓ is even, take $i = \frac{\ell}{2}$. It is easy to check that $P_{\frac{\ell}{2}} = (2,1,4,3,6,5,\ldots)$. By statement (e), if $\lceil \frac{n}{2} \rceil + 2 \geq 6$, then $c(1,5) = c(5,8) \neq c(5,6)$, a contradiction. So $n = 6$. Since $3 \in [1, y_s]$, it yields that $c(3,4) = c(3,6)$ by Claim 2 (a); by Claim 2 (b), $c(1,3) = c(3,4)$. Therefore, $c(1,3) = c(3,4) = c(3,6)$, which contradicts the fact that $\Delta^{\text{mon}}(K_6) < \lfloor \frac{6}{2} \rfloor$. $\square$

Claim 7. $c(y_1+1, y_1+3) \notin \{c(y_1+1, y_1+2), c(y_1+3, y_1+4)\}$.

Proof of Claim 7. By Claim 3, $S(P^*) = [t,u]$ and $y_1 + 3 \in N^c(y_1+1, P^*)$, so $c(y_1+1, y_1+2) \neq c(y_1+1, y_1+3)$. For the longest properly colored path $Q^* = (y_1+3, y_1+4, \ldots, \ell, y_1+2, y_1+1, \ldots, 1) = (\psi(1), \psi(2), \ldots, \psi(\ell))$, we know that $S(Q^*) = [t,u]$ and $N^c(y_1+3, Q^*) = \{\psi(j) \mid j \in A\}$, where

$$A = \begin{cases} [3, y_1''] \cup [t'', u''] \cup [w'', \ell] & \text{if } t > y_1 + 3; \quad (2.6) \\ [t'', u''] \cup [w'', \ell-1] & \text{if } t = y_1 + 3, \quad (2.7) \end{cases}$$

for some $2 \leq y_1'' < t'' < u'' < w'' \leq \ell$. Since $y_1+2 \in S(P_0)$, $c(y_1+2, y_1+3) = c(y_1+2, \ell)$. Thus $y_1+2 \in \{\psi(j) \mid j \in [w'', \ell-1]\}$, so $[2, y_1+2] \subseteq \{\psi(j) \mid j \in [w'', \ell-1]\}$. In particular, $y_1+1 \in N^c(y_1+3, Q^*)$, so $c(y_1+1, y_1+3) \neq c(y_1+3, y_1+4)$, which completes the proof of Claim 7. $\square$



Fig. 2: Cycle $(y_1+1, y_1+2, \ell, \ell-1, \ldots, y_1+3, y_1+1)$

By Claim 7, note that $(y_1+1, y_1+2, \ell, \ell-1, \ldots, y_1+3, y_1+1)$ is a properly colored cycle (see Fig. 2) containing $Y \cup \{\ell-1, \ell\} \setminus \{y_1\}$, then its length is exactly $\lceil \frac{n}{2} \rceil + 1$. Moreover, $Y = [\ell - \lceil \frac{n}{2} \rceil - 1, \ell - 2]$.

If $\ell = n$, then $Y = [\lfloor \frac{n}{2} \rfloor - 1, n-2]$ and $y_1 = \lfloor \frac{n}{2} \rfloor - 1$. By Claim 3 and Claim 6, $u \geq y_s + 3$. Now

$$\left\lceil \frac{n}{2} \right\rceil + 1 \geq u \geq y_s + 3 = y_1 + |S(P_0)| + 2 \geq \left\lfloor \frac{n}{2} \right\rfloor + 4, \tag{2.8}$$

a contradiction.

Then we may assume that $\ell < n$, i.e., there exists a vertex z which is not on P_0. We know that $c(\ell - 1, \ell) = c(\ell, z)$, otherwise $(1, 2, \ldots, \ell, z)$ is a properly colored path which is longer than P_0. We have known that $(1, 2, \ldots, y_1 + 1, \ell, \ell - 1, \ldots, y_1 + 2)$ is a longest properly colored path, it yields that $c(y_1 + 2, y_1 + 3) = c(y_1 + 2, z)$ for the same reason. Since $y_2 + 2 \in S(P_0)$, we have $c(y_1 + 2, y_1 + 3) = c(y_1 + 2, \ell) \neq c(\ell - 1, \ell) = c(\ell, z)$, and hence $c(y_1 + 2, z) \neq c(\ell, z)$. Therefore, $(1, \ldots, y_1, \ell, z, y_1 + 2, y_1 + 1, y_1 + 3, \ldots, \ell - 1)$ is a properly colored path which is longer than P_0, a contradiction.

References

[1] N. Alon and G. Gutin, Properly colored Hamilton cycles in edge-colored complete graphs, Random Structures and Algorithms 11 (1997) (2) 179–186.

[2] J. Bang-Jensen and G. Gutin, Alternating cycles and paths in edge-coloured multigraphs: A survey, Discrete Math. 165-166 (1997) 39–60.

[3] J. Bang-Jensen, G. Gutin and A. Yeo, Properly coloured Hamiltonian paths in edge-coloured complete graphs, Discrete Appl. Math. 82 (1998) (1-3) 247–250.

[4] O. Barr, Properly coloured Hamiltonian paths in edge-coloured complete graphs without monochromatic triangles, Ars Combin. 50 (1998) 316–318.

[5] B. Bollobás and P. Erdös, Alternating hamiltonian cycles, Israel J. Math. 23 (1976) (2) 126–131.

[6] C. C. Chen and D. E. Daykin, Graphs with Hamiltonian cycles having adjacent lines different colors, J. Combin. Theory Ser. B 21 (1976) (2) 135–139.

[7] J. Feng, H.-E. Giesen, Y. Guo, G. Gutin, T. Jensen and A. Rafiey, Characterization of edge-colored complete graphs with properly colored Hamilton paths, J. Graph Theory 53 (2006) (4) 333–346.

[8] S. Fujita and C. Magnant, Properly colored paths and cycles, Discrete Appl. Math. 159 (2011) (14) 1391–1397.

[9] J. W. Grossman and R. Häggkvist, Alternating cycles in edge-partitioned graphs, J. Combin. Theory Ser. B 34 (1983) (1) 77–81.

[10] G. Gutin and E. J. Kim, Properly coloured cycles and paths: Results and open problems, Lecture Notes in Comput. Sci. 5420 (2009) 200–208.

[11] M. Kano and X. Li, Monochromatic and heterochromatic subgraphs in edge-colored graphs - a survey, Graphs Combin. 24 (2008) (4) 237–263.

[12] H. Li, G. Wang and S. Zhou, Long alternating cycles in edge-colored complete graphs, Lecture Notes in Comput. Sci. 4613 (2007) 305–309.

[13] A. S. L. Lo, A Dirac type condition for properly coloured paths and cycles, arXiv:1008.3242.

[14] L. Pósa, Hamiltonian circuits in random graphs, Discrete Math. 14 (1976) (4) 359–364.

[15] J. Shearer, A property of the colored complete graph, Discrete Math. 25 (1979) (2) 175–178.

[16] A. Yeo, A note on alternating cycles in edge-coloured graphs, J. Combin. Theory Ser. B 69 (1997) (2) 222–225.